arXiv:hep-th/9403041v1 7 Mar 1994

THEF-NYM 94.2
hep-th/9403041
March 1994

EXACT STRING BACKGROUND FROM A WZW MODEL BASED ON THE HEISENBERG GROUP

Alexandros A. Kehagias *†
and
Patrick A. A. Meessen ‡§

Institute of Theoretical Physics
University of Nijmegen, Toernooiveld 1,
6525 ED, Nijmegen, The Netherlands

Abstract

A WZW model for the non-semi-simple D-dimensional Heisenberg group, which directly generalizes the E_2^c group, is constructed. It is found to correspond to an D-dimensional string background of plane-wave type with physical Lorentz signature D-2. Perturbative and non-perturbative considerations lead both to an integer central charge c=D.

---

[*] Partially supported by CEC Contract No. ERBCHBGCT920197

[†] e-mail: kehagias@sci.kun.nl

[‡] Partially supported by 'de Wet op Studiefinanciering' No. 5614-71105-0-13

[§] e-mail: patrickm@sci.kun.nl

Recently, WZW models based on non-semi-simple groups have attained a lot of interest [1–6], particularly because they lead to exactly solvable models corresponding to exact string backgrounds. To write down a WZW model for a non-semi-simple group is not straightforward since its Killing metric is degenerate. However, it is possible that another, non-degenerate, invariant bilinear form on the algebra exists which allows for a WZW model to be constructed. The non-semi-simple groups explicitly studied so far are the centrally extended 2D Euclidean group E_2^c [1], its d-dimensional generalization E_d^c [3], as well as a large class of non-semi-simple groups arising after contraction of G×H where H is a subgroup of G [2]. In particular, E_2^c is isomorphic to the 4-dimensional Heisenberg group. It is then natural to discuss its D-dimensional analogue, namely the Heisenberg group H_D. For this, a non-degenerate bilinear form exists, a WZW model can be written down and a Sugawara-type construction can be carried out. The resulting D-dimensional conformal σ-model has one null Killing vector and the corresponding conformal field theory has $c = D$, in accordance with the general result of [4].

The WZW models based on a group G are defined on a two-surface Σ by the action

$$\mathcal{S}_{WZW} \;=\; -\frac{k}{4\pi}\int_\Sigma Tr(g^{-1}dg\, g^{-1}dg) + \frac{k}{6\pi}\int_B Tr(g^{-1}dg\wedge g^{-1}dg\wedge g^{-1}dg)\,, \qquad (1)$$

where $g^{-1}dg$ are the right invariant 1-forms on G and B is a three manifold bounded by Σ. The right invariant 1-forms are elements of the Lie algebra of G, so that they may be expressed as

$$g^{-1}dg \;=\; iA^I T_I\,, \qquad (2)$$

where T_I $(I = 1,\ldots,dimG)$ are the generators of the group satisfying

$$[T_I, T_J] \;=\; if_{IJ}{}^K T_K\,. \qquad (3)$$

The WZW action (1) can then be written in terms of the A^I's as

$$\mathcal{S}_{WZW} \;=\; -\frac{k}{4\pi}\int_\Sigma d^2\sigma A^I_\alpha A^{J\alpha}\Omega_{IJ} \;-\; \frac{k}{12\pi}\int_B d^3\sigma\, \epsilon^{\alpha\beta\gamma} A^I_\alpha A^J_\beta A^L_\gamma f_{IJ}{}^N\Omega_{NL}\,, \qquad (4)$$

where $\Omega_{IJ} = Tr(T_I T_J)$ is the Killing metric for the group G. However, when the group is non-semi-simple, the Killing metric is degenerate and one might replace it, as was proposed in [1], by a symmetric, invariant and non-degenerate form Ω_{IJ} that satisfies

$$f_{IJ}{}^K\Omega_{KL} \;+\; f_{IL}{}^K\Omega_{KJ} \;=\; 0\,. \qquad (5)$$

Such a form can be found, apart from the above mentioned cases, also for the Heisenberg group H_D.

Before proceeding to the general case, let us first examine the case H$_4$, which originates from the dynamics of a single one-dimensional harmonic oscillator. H$_4$ is generated by $\{\alpha,\, \alpha^\dagger,\, N=\alpha^\dagger\alpha,\, I\}$ and the commutation relations

$$\begin{aligned} [\alpha,\alpha^\dagger] &= I\,, \\ [N,\alpha^\dagger] &= \alpha^\dagger\,, \\ [N,\alpha] &= -\alpha\,. \end{aligned} \tag{6}$$

It is non-semi-simple so that its Killing form is degenerate. A non-degenerate solution to eq.(5) exists and is given by

$$\Omega_{IJ} \;=\; \begin{pmatrix} 0 & a & 0 & 0 \\ a & 0 & 0 & 0 \\ 0 & 0 & b & -a \\ 0 & 0 & -a & 0 \end{pmatrix}\,,\; \Omega^{IJ} \;=\; \begin{pmatrix} 0 & \frac{1}{a} & 0 & 0 \\ \frac{1}{a} & 0 & 0 & 0 \\ 0 & 0 & 0 & -\frac{1}{a} \\ 0 & 0 & -\frac{1}{a} & -\frac{b}{a^2} \end{pmatrix}\,. \tag{7}$$

A general element of H$_4$ is written as

$$g \;=\; e^{iq\alpha+i\bar{q}\alpha^\dagger}e^{iuN+ivI}\,, \tag{8}$$

where the first term on the right hand side of eq.(8) is an element of the coset space H$_4/U(1)\times U(1) \sim \mathbb{C}$ parametrized by the complex coordinates $q,\bar{q}$. By using the relations

$$\begin{aligned} e^{iq\alpha+i\bar{q}\alpha^\dagger} &= e^{iq\alpha}e^{i\bar{q}\alpha^\dagger}e^{\frac{q\bar{q}}{2}}\,, \\ e^{-iuN}\alpha e^{iuN} &= e^{iu}\alpha\,, \\ e^{-i\bar{q}\alpha^\dagger}\alpha e^{i\bar{q}\alpha^\dagger} &= \alpha\,+\,i\bar{q}I\,, \end{aligned} \tag{9}$$

we find that

$$g^{-1}dg \;=\; ie^{iu}dq\alpha\,+\,ie^{-iu}d\bar{q}\alpha^\dagger\,+\,iduN\,+\,(idv+\frac{1}{2}qd\bar{q}-\frac{1}{2}\bar{q}dq)I\,, \tag{10}$$

so that the A^I's in eq.(2) are given by

$$\begin{aligned} A^1 &= e^{iu}dq\,, \\ A^2 &= e^{-iu}d\bar{q}\,, \\ A^3 &= du\,, \\ A^4 &= dv\,-\,\frac{i}{2}qd\bar{q}\,+\,\frac{i}{2}\bar{q}dq\,. \end{aligned} \tag{11}$$

The terms that are being integrated over in (4) are calculated to be

$$\begin{aligned} A^I_\alpha A^{J\alpha}\Omega_{IJ} &= 2a\partial_\alpha q\partial^\alpha\bar{q} - 2a(\partial_\alpha v - \frac{i}{2}q\partial_\alpha\bar{q} + \frac{i}{2}\bar{q}\partial_\alpha q)\partial^\alpha u + b\partial_\alpha u\partial^\alpha u\,, \\ \epsilon^{\alpha\beta\gamma}A^I_\alpha A^J_\beta A^K_\gamma f_{IJK} &= 6ia\epsilon_{\alpha\beta\gamma}\partial^\gamma(u\partial^\alpha q\partial^\beta\bar{q})\,, \end{aligned} \tag{12}$$

and the WZW action (4) is written as

$$\begin{aligned}\mathcal{S}_{WZW} &= -\frac{k}{4\pi}\int d^2\sigma\Big(2a\partial_\alpha q\partial^\alpha\bar{q} - 2a(\partial_\alpha v - \frac{i}{2}q\partial_\alpha\bar{q} + \frac{i}{2}\bar{q}\partial_\alpha q)\partial^\alpha u \\ &\quad + b\partial_\alpha u\partial^\alpha u + 2ia\epsilon_{\alpha\beta}u\partial^\alpha q\partial^\beta\bar{q}\Big)\,. \end{aligned} \tag{13}$$

By regarding this action as a σ-model action of the form

$$\mathcal{S} \;=\; \int d^2\sigma(G_{\mu\nu}\partial_\alpha X^\mu\partial^\alpha X^\nu \;+\; B_{\mu\nu}\epsilon^{\alpha\beta}\partial_\alpha X^\mu\partial_\beta X^\nu)\,, \tag{14}$$

we can read off the background space-time metric and the antisymmetric field. In the coordinate base $[dq, d\bar{q}, du, dv]$ they are given, up to multiplicative factors, by

$$\begin{aligned} G_{\mu\nu} &= \begin{pmatrix} 0 & \frac{1}{2} & -\frac{i}{4}\bar{q} & 0 \\ \frac{1}{2} & 0 & \frac{i}{4}q & 0 \\ -\frac{i}{4}\bar{q} & \frac{i}{4}q & \beta^2 & \frac{1}{2} \\ 0 & 0 & \frac{1}{2} & 0 \end{pmatrix}, \\ B_{q\bar{q}} &= \frac{i}{2}u\,, \end{aligned} \tag{15}$$

where $\beta^2 = \frac{b}{2a}$, and thus, the background space-time line element is given by

$$ds^2 \;=\; dqd\bar{q} - (dv - \frac{i}{2}qd\bar{q} + \frac{i}{2}\bar{q}dq)du + \beta^2 du^2\,. \tag{16}$$

It describes a plane-wave space-time [7, 8] and it can be identified to the one found by Nappi and Witten in [1] by virtue of the transformation $q = a_1 - ia_2$ and $\bar{q} = a_1 + ia_2$. By introducing polar coordinates $q = Re^{i\theta}$, $\bar{q} = Re^{-i\theta}$, the line element turns out to be

$$ds^2 \;=\; dR^2 + R^2d\theta^2 - (dv - R^2d\theta)du + \beta^2du^2\,. \tag{17}$$

The signature of this metric is manifest in the orthonormal base

$$\begin{aligned} e^0 &= \frac{1}{2\beta}(dv - R^2d\theta)\,, \\ e^1 &= dR\,, \\ e^2 &= Rd\theta\,, \\ e^3 &= \beta du - e^0\,, \end{aligned} \tag{18}$$

where the metric is $\eta_{\mu\nu} = (-1, +1, +1, +1)$. The only non-vanishing components of the Ricci tensor in the above base are

$$R_{00} \;=\; R_{33} \;=\; R_{03} \;=\; \frac{1}{2\beta^2}\,. \tag{19}$$

In the same base, the antisymmetric two form field $B = \frac{1}{2} B_{\mu\nu} dx^\mu \wedge dx^\nu$ is written as

$$B \; = \; u e^1 \wedge e^2 \,, \tag{20}$$

and thus

$$H \; = \; dB \; = \; \frac{1}{\beta} \, e^3 \wedge e^1 \wedge e^2 + \frac{1}{\beta} \, e^0 \wedge e^1 \wedge e^2 \,. \tag{21}$$

The non-vanishing components of H are then

$$H_{123} \; = \; H_{012} \; = \; \frac{1}{\beta} \,. \tag{22}$$

Employing eqs.(19), (22) in the one-loop beta-function equations

$$\begin{aligned}
0 \; &= \; -R + \frac{1}{12} H^2 + 4(\nabla\Phi)^2 - 4\nabla^2\Phi + \frac{2(c-4)}{3\alpha'} \,, \\
0 \; &= \; R_{\mu\nu} - \frac{1}{4} H^2_{\mu\nu} + 2\nabla_\mu \nabla_\nu \Phi \,, \\
0 \; &= \; \nabla_\lambda H^\lambda_{\mu\nu} - 2(\nabla_\lambda \Phi) H^\lambda_{\mu\nu} \,,
\end{aligned} \tag{23}$$

one can find that the dilaton is constant and that the central charge is four ($c = 4$). (A constant dilaton is expected on behalf of the homogeneity of the space.)

Let us now discuss the general case H_D. The Heisenberg group H_D is a D-dimensional group generated by the set of generators $\{\alpha_i, \, \alpha_i^\dagger, \, N = \sum_i \alpha_i^\dagger \alpha_i, \, I, \, i = 1, \ldots, r = \frac{D-2}{2}\}$ and the commutation relations

$$\begin{aligned}
[\alpha_i, \alpha_j^\dagger] \; &= \; \delta_{ij} I \,, \\
[N, \alpha_i] \; &= \; -\alpha_i \,, \\
[N, \alpha_i^\dagger] \; &= \; \alpha_i^\dagger \,, \\
[N, I] \; &= \; [\alpha_i, I] \; = \; [\alpha_i^\dagger, I] \; = \; 0 \,.
\end{aligned} \tag{24}$$

The Killing form of H_D is degenerate and a non-degenerate solution to eq.(5) which can replace the Killing form is given by

$$\Omega_{IJ} \; = \; \begin{pmatrix} A_1 & & & & \\ & \ddots & & & \\ & & A_r & & \\ & & & b & -a \\ & & & -a & 0 \end{pmatrix} \,, \tag{25}$$

where

$$A_1 \; = \; A_2 \; = \ldots = \; A_r \; = \; \begin{pmatrix} 0 & a \\ a & 0 \end{pmatrix} \,. \tag{26}$$

An element of H_D is expressed as

$$g \;=\; e^{i\sum_i (q_i\alpha_i + \bar{q}_i\alpha_i^\dagger)} e^{iuN+ivI}\,, \tag{27}$$

and proceeding as before, we find that

$$\begin{aligned}
A^i &= e^{iu}dq_i\,,\\
\bar{A}^i &= e^{-iu}d\bar{q}_i\,,\\
A^{D-1} &= du\,,\\
A^D &= dv - \frac{i}{2}\sum_i (q_i d\bar{q}_i - \bar{q}_i dq_i)\,.
\end{aligned} \tag{28}$$

The WZW action is then written as

$$\begin{aligned}
\mathcal{S}_{WZW} \;=\; & -\frac{k}{4\pi}\int_\Sigma d^2\sigma \left(2a\sum_i \partial_\alpha q_i \partial^\alpha \bar{q}_i - 2a(\partial_\alpha v - \frac{i}{2}\sum_i q_i\partial_\alpha\bar{q}_i - \bar{q}_i\partial_\alpha q_i)\partial^\alpha u \right.\\
+\;& \left. b\partial_\alpha u \partial^\alpha u + 2ia\epsilon_{\alpha\beta} u \sum_i \partial^\alpha q_i \partial^\beta \bar{q}_i \right)\,.
\end{aligned} \tag{29}$$

By comparing this action with eq.(14) we can read off the space-time metric, which is written, up to multiplicative factors, as

$$G_{\mu\nu} \;=\; \begin{pmatrix}
0 & \frac{1}{2} & & & & -\frac{i}{4}\bar{q}_1 & 0\\
\frac{1}{2} & 0 & & & & \frac{i}{4}q_1 & 0\\
 & & \ddots & & & \vdots & \vdots\\
 & & & 0 & \frac{1}{2} & -\frac{i}{4}\bar{q}_r & 0\\
 & & & \frac{1}{2} & 0 & \frac{i}{4}q_r & 0\\
-\frac{i}{4}\bar{q}_1 & \frac{i}{4}q_1 & \dots & -\frac{i}{4}\bar{q}_r & \frac{i}{4}q_r & \beta^2 & \frac{1}{2}\\
0 & 0 & \dots & 0 & 0 & \frac{1}{2} & 0
\end{pmatrix}, \tag{30}$$

$$G^{\mu\nu} \;=\; \begin{pmatrix}
0 & 2 & & & & & -iq_1\\
2 & 0 & & & & & i\bar{q}_1\\
 & & \ddots & & & & \vdots\\
 & & & 0 & 2 & & -iq_r\\
 & & & 2 & 0 & & i\bar{q}_r\\
 & & & & & 0 & 2\\
-iq_1 & i\bar{q}_1 & \dots & -iq_r & i\bar{q}_r & 2 & \gamma - 4\beta^2
\end{pmatrix}, \tag{31}$$

where the notation $\gamma = \sum_i q_i\bar{q}_i$ has been used. The antisymmetric field then is found to be

$$B_{q_i\bar{q}_j} \;=\; \frac{i}{2}u\delta_{ij}\,. \tag{32}$$

The line element for this case is then written as

$$ds^2 \;=\; \sum_i dq_i d\bar{q}_i - \left(dv - \frac{i}{2}\sum_i (q_i d\bar{q}_i - \bar{q}_i dq_i)\right) du + \beta^2 du^2 \,. \tag{33}$$

Introducing polar coordinates $q_i = R_i e^{i\theta_i}$, $\bar{q}_i = R_i e^{-i\theta_i}$ we find that the metric is $\eta_{\mu\nu} = (-1, +1, \ldots, +1)$ in the base

$$\begin{aligned} e^0 &= \frac{1}{2\beta}(dv - \sum_i R_i^2 d\theta_i)\,, \\ e^{2i-1} &= dR_i\,, \\ e^{2i} &= R_i d\theta_i\,, \\ e^{D-1} &= \beta du - e^0\,, \end{aligned} \tag{34}$$

where $i = 1, \ldots, r$. The non-vanishing components of the Ricci tensor in the base (34) are then

$$R^{D-1}{}_{D-1} \;=\; -R^0{}_0 \;=\; -R^0{}_{D-1} \;=\; \frac{D-2}{4\beta^2}\,, \tag{35}$$

whereas the antisymmetric 3-form field is given by

$$H \;=\; dB \;=\; \frac{1}{\beta}\sum_{i=1}^{r}\left(e^{D-1}\wedge e^{2i-1}\wedge e^{2i} \;+\; e^0 \wedge e^{2i-1}\wedge e^{2i}\right)\,. \tag{36}$$

It is straightforward then to verify that the vanishing of the one-loop beta functions in eqs.(23) gives $c = D$.

One can understand this result perturbatively by considering the loop diagrams that contribute to the beta function. Since the propagator of the u field vanishes due to $G^{uu} = G^{D-1\,D-1} = 0$, the only diagrams that can be drawn are one-loop diagrams with two external u lines and q_i, $\bar{q}_i$ in the internal lines. However, the interaction terms in (29) give opposite contributions to these diagrams, in exactly the same way discussed by Nappi and Witten [1, 9] leading to $c = D$ identically in perturbation theory.

The same result can be obtained non-perturbatively as follows [4, 5, 10]: The Noether currents associated with the WZW model (4) have the following OPE

$$J_I(z)J_J(w) \;=\; -\frac{k\,\Omega_{IJ}}{2(z-w)^2} + f_{IJ}{}^K \frac{J_K(w)}{(z-w)} + reg. \tag{37}$$

Let us define the bilinear in the currents energy-momentum tensor

$$T(z) \;=\; L^{IJ}\; {:}J_I J_J{:}(z)\,. \tag{38}$$

The condition for the J_I's to be primary fields of weight 1, is written as

$$T(z)J_I(w) \;=\; \frac{J_I(w)}{(z-w)^2} + \frac{\partial J_I(w)}{(z-w)} + reg. \tag{39}$$

and by using eq.(38) we get [4, 5]

$$\begin{aligned} L^{IJ} f_{KJ}{}^{L} + L^{JL} f_{KJ}{}^{I} &= 0 \,, && (40) \\ 2L^{IJ}\Omega_{KJ} + L^{MN} f_{KM}{}^{L} f_{LN}{}^{I} &= \delta^{I}{}_{K} \,. && (41) \end{aligned}$$

Thus L^{IJ} is an invariant symmetric tensor as follows from eq.(40) and employing the latter in eq.(41) we get

$$L^{IJ}(2\Omega_{KJ} + \kappa_{KJ}) = \delta^{I}{}_{K} \,, \tag{42}$$

where $\kappa_{KJ} = f_{KM}{}^{L} f_{JL}{}^{M}$ is the Killing form. Then it follows that L^{IJ} is the inverse of the matrix $(2\Omega_{KJ} + \kappa_{KJ})$ and in our case one finds that

$$L^{IJ} = \frac{1}{2a} \begin{pmatrix} 0 & 1 & & & & & \\ 1 & 0 & & & & & \\ & & \ddots & & & & \\ & & & 0 & 1 & & \\ & & & 1 & 0 & & \\ & & & & & 0 & -1 \\ & & & & & -1 & -\frac{b+r}{a} \end{pmatrix} . \tag{43}$$

The central charge is given by

$$c = 2L^{IJ}\Omega_{IJ} = D \,, \tag{44}$$

in accordance with the results in [4] since one may look upon the Heisenberg algebra H_D as the double $U(1)$ extention of the algebra

$$[\alpha_i, \alpha_j{}^{\dagger}] = 0 \,. \tag{45}$$

The above construction is a direct generalization of Nappi and Witten's result [1], which is recovered in the case $D = 4$. The space-time described by the metric (33) is homogeneous, it has $2(D-1)$ space-like Killing vectors and a null one, a total of $2D-1$. It describes a plane-wave type space-time with flat wave surfaces ($u, v = const.$) since the coset space $\mathrm{H}_D/U(1)\times U(1)$ is isomorphic to $\mathbb{C}^{r}$. It has one time-like direction (Lorentz signature $D-2$) and thus it can be considered as a D-dimensional string background which can replace Minkowski space-time.

We would like to thank P. de Boer for his help in the computer programm and N. Obers for discussions.

References

[1] C.R. Nappi and E. Witten, Phys. Rev. Lett. 71 (1993)3751.

[2] D.I. Olive, E. Rabinovici and A. Schwimmer, Phys. Lett. B 321 (1994)361.

[3] K. Sfetsos, *"Exact string backgrounds from WZW models based on non-semi-simple groups"*, preprint THU-93/31, hep-th/9311093; *"Gauged WZW models and non-abelian duality"*, preprint THU-94/1, hep-th/9402031.

[4] J.M. Figueroa-O'Farrill and S. Stanciu, *"Nonsemisimple Sugawara constructions"*, preprint QMW-PH-94-2, hep-th/9402035.

[5] N. Mohammedi, *"On bosonic and supersymmetric current algebras for non-semi-simple groups"*, preprint BONN-HE-93-51, hep-th/9312182.

[6] E. Kiritsis and C. Kounnas, Phys. Lett. B 320 (1994)264.

[7] H.W. Brinkmann, Proc. Nat. Acad. Sci. (U.S.) 9 (1923) 1;
N. Rosen, Phys. Z. Sowjetunion, 12.4 (1937)366;
J. Ehlers and W. Kundt, in *Gravitation: an introduction to current research*, ed. L. Witten, J.W., N.Y. 1962.

[8] R. Güven, Phys. Lett. B 191 (1987)275;
D. Amati and C. Klimčik, Phys. Lett. B 219 (1989)443;
G.T. Horowitz and A.R. Steif, Phys. Rev. Lett. 64 (1990)260;
A.A. Tseytlin, Nucl. Phys. B 390 (1993)153; Phys. Rev. D 47 (1993)3421.

[9] C.R. Nappi, Phys. Rev. D 21 (1980)418;
E. Witten, Commun. Math. Phys. 92 (1984)455.

[10] M.B. Halpern and E. Kiritsis, Mod. Phys. Lett. A 4 (1989)1373; Erratum Mod. Phys. Lett. A 4 (1989)1797.